

D8S1132

Population	Ref	n	15	16	17	18	19	20	21	22	23	24	25
Europe													
Germany (pooled)		815	0.0006	0.0141	0.1098	0.1914	0.1348	0.1528	0.1282	0.1234	0.1001	0.0348	0.0098
Germany (Düsseldorf)	3	484	0.0010	0.0176	0.1167	0.1736	0.1291	0.1632	0.1250	0.1271	0.1023	0.0351	0.0093
Germany (Halle)	1, 2	100	0.0000	0.0100	0.0850	0.2300	0.1050	0.1600	0.1400	0.1450	0.0700	0.0400	0.0150
Germany (Münster)	1	134	0.0000	0.0040	0.0930	0.1940	0.1450	0.1310	0.1420	0.1120	0.1420	0.0220	0.0150
Germany (Wiesbaden)	1, 2	97	0.0000	0.0150	0.1240	0.2370	0.1800	0.1240	0.1130	0.0980	0.0620	0.0460	0.0000
Asia, Far East													
China (Shenyang)	4	140	0.0000	0.0300	0.0820	0.1940	0.1980	0.1490	0.1160	0.1270	0.0820	0.0190	0.0040
Japan (Sendai)	4	120	0.0000	0.0000	0.0960	0.1830	0.1460	0.1380	0.1330	0.1460	0.1170	0.0380	0.0040

References

- (1) Wiegand, P., Schneider, H.R., Schürenkamp, M., Kleiber, M., Brinkmann, B. (1998) Tetranucleotide STR system D8S1132: sequencing data and population genetic comparisons. *Int. J. Legal Med.* 111: 180-182
- (2) Schneider, H.R., Wiegand, P., Schürenkamp, M., Kleiber, M. (1998) STR system D8S1132: Genetic data of two German population samples. *Progr. Forensic Genet.* 7: 350-352
- (3) Huckenbeck, W., Scheil, H.-G. (2001) unpublished data
- (4) He, P., Nata, M., Kanetake, J., Ji, G., Yan, W., Li, X., Sagisaka, K. (1997) Polymorphism of short tandem repeat (STR) loci PLA2, D3S2459, D8S315 (kw38), CYP19, D3S1359, FGA, and D8S1132 in the Japanese and Chinese populations. *Tohoku J. Exp. Med.* 183, 251-262